
Vocabulary Flash Cards	
acute angle	adjacent angles
Chapter 1 (p. 39)	Chapter 1 (p. 48)
angle	angle bisector
Chapter 1 (p .38)	Chapter 1 (p. 42)
axiom	between
Chapter 1 (p. 12)	Chapter 1 (p. 14)
collinear points	complementary angles
Chapter 1 (p. 4)	Chapter 1 (p. 48)

Two angles that share a common vertex and side, but have no common interior points




 $\angle 5$ and $\angle 6$ are adjacent angles.

An angle that has a measure greater than 0° and less than 90°




A ray that divides an angle into two angles that are congruent



 \overrightarrow{YW} bisects $\angle XYZ$, so $\angle XYW \cong \angle ZYW$.

A set of points consisting of two different rays that have the same endpoint



When three points are collinear, one point is between the other two.




Point B is between points A and C.

A rule that is accepted without proof

The Segment Addition Postulate states that if B is between A and C, then AB + BC = AC.

Two angles whose measures have a sum of 90°



 $\angle BAC$ and $\angle CAB$ are complementary angles.


Points that lie on the same line



A, B, and C are collinear.


Vocabulary Flash Cards	
congruent angles	congruent segments
Chapter 1 (p. 40)	Chapter 1 (p. 13)
construction	coordinate Chapter 1 (p. 12)
Chapter 1 (p. 13)	Chapter 1 (p. 12)
coplanar points	defined terms
Chapter 1 (p. 4)	Chapter 1 (p. 5)
distance	endpoints
Chapter 1 (p. 12)	Chapter 1 (p. 5)

Line segments that have the same length




 $\overline{AB} \cong \overline{CD}$


Two angles that have the same measure



A real number that corresponds to a point on a line




A geometric drawing that uses a limited set of tools, usually a compass and a straightedge



Terms that can be described using known words, such as *point* or *line*


Line segment and ray are two defined terms.

Points that lie in the same plane



A, B, and C are coplanar.

Points that represent the ends of a line segment or ray



The absolute value of the difference of two coordinates on a line

$$A \qquad AB \qquad B$$

$$x_1 \qquad x_2$$

$$AB = |x_2 - x_1|$$

Vocabulary Flash Cards exterior of an angle interior of an angle Chapter 1 (p. 38) Chapter 1 (p. 38) intersection line *Chapter 1 (p. 4) Chapter 1 (p. 6)* line segment linear pair Chapter 1 (p. 50) *Chapter 1 (p. 5)* measure of an angle midpoint Chapter 1 (p. 20) *Chapter 1 (p. 39)*


The region that contains all the points between the sides of an angle




The region that contains all the points outside of an angle



A line has one dimension. It is represented by a line with two arrowheads, but it extends without end.




The set of points two or more geometric figures have in common



The intersection of two different lines is a point.

Two adjacent angles whose noncommon sides are opposite rays




Consists of two endpoints and all the points between them



The point that divides a segment into two congruent segments



M is the midpoint of AB. So, $\overline{AM}\cong\overline{MB}$ and AM=MB. The absolute value of the difference between the real numbers matched with the two rays that form the angle on a protractor




Big Ideas Math Geometry

Vocabulary Flash Cards obtuse angle opposite rays *Chapter 1 (p. 5)* Chapter 1 (p. 39) plane point *Chapter 1 (p. 4) Chapter 1 (p. 4)* postulate ray *Chapter 1 (p. 5) Chapter 1 (p. 12)* right angle segment *Chapter 1 (p. 5) Chapter 1 (p. 39)*


If point C lies on \overrightarrow{AB} between A and B, then \overrightarrow{CA} An angle that has a measure greater than 90° and and \overline{CB} are opposite rays. less than 180° \overrightarrow{CA} and \overrightarrow{CB} are opposite rays. A location in space that is represented by a dot and A flat surface made up of points that has two has no dimension dimensions and extends without end, and is represented by a shape that looks like a floor or a wall point A plane M, or plane ABC \overrightarrow{AB} is a ray if it consists of the endpoint A and all A rule that is accepted without proof points on \overrightarrow{AB} that lie on the same side of A as B. The Segment Addition Postulate states that if *B* is between A and C, then AB + BC = AC. \overrightarrow{AB} Consists of two endpoints and all the points An angle that has a measure of 90° between them

Vocabulary Flash Cards	
segment bisector	sides of an angle
Chapter 1 (p. 20)	Chapter 1 (p. 38)
straight angle	supplementary angles
Chapter 1 (p. 39)	Chapter 1 (p. 48)
undefined terms	vertex of an angle
Chapter 1 (p. 4)	Chapter 1 (p. 38)
vertical angles Chapter 1 (p. 50)	

The rays of an angle




A point, ray, line, line segment, or plane that intersects the segment at its midpoint



 \overrightarrow{CD} is a segment bisector of \overrightarrow{AB} . So, $\overrightarrow{AM} \cong \overrightarrow{MB}$ and $\overrightarrow{AM} = \overrightarrow{MB}$.

Two angles whose measures have a sum of 180°




An angle that has a measure of 180°



 $\angle JKM$ and $\angle LKM$ are supplementary angles.


The common endpoint of the two rays that form an angle



Words that do not have formal definitions, but there is agreement about what they mean

In geometry, the words *point*, *line*, and *plane* are undefined terms.

Two angles whose sides form two pairs of opposite rays



 $\angle 3$ and $\angle 6$ are vertical angles.

 $\angle 4$ and $\angle 5$ are vertical angles.